

language: English

12 July 2006

Problem 1. Let ABC be a triangle with incentre I. A point P in the interior of the triangle satisfies

$$\angle PBA + \angle PCA = \angle PBC + \angle PCB$$
.

Show that $AP \geq AI$, and that equality holds if and only if P = I.

Problem 2. Let P be a regular 2006-gon. A diagonal of P is called good if its endpoints divide the boundary of P into two parts, each composed of an odd number of sides of P. The sides of P are also called good.

Suppose P has been dissected into triangles by 2003 diagonals, no two of which have a common point in the interior of P. Find the maximum number of isosceles triangles having two good sides that could appear in such a configuration.

Problem 3. Determine the least real number M such that the inequality

$$\left| ab(a^2 - b^2) + bc(b^2 - c^2) + ca(c^2 - a^2) \right| \le M(a^2 + b^2 + c^2)^2$$

holds for all real numbers a, b and c.



language: English

13 July 2006

Problem 4. Determine all pairs (x, y) of integers such that

$$1 + 2^x + 2^{2x+1} = y^2.$$

Problem 5. Let P(x) be a polynomial of degree n > 1 with integer coefficients and let k be a positive integer. Consider the polynomial $Q(x) = P(P(\ldots P(P(x)) \ldots))$, where P occurs k times. Prove that there are at most n integers t such that Q(t) = t.

Problem 6. Assign to each side b of a convex polygon P the maximum area of a triangle that has b as a side and is contained in P. Show that the sum of the areas assigned to the sides of P is at least twice the area of P.